

# STELLAR POPULATIONS AND THE LOCAL GROUP MEMBERSHIP OF THE DWARF GALAXY DDO 210<sup>1</sup>

MYUNG GYOON LEE

Department of Astronomy, Seoul National University, Seoul 151-742, Korea

Electronic mail: mglee@astro.snu.ac.kr

ANTONIO APARICIO

Instituto de Astrofísica de Canarias, La Laguna, Tenerife, Canary Islands, Spain

Electronic mail: aaj@iac.es

NIKOLAY TIKONOV

Special Astrophysical Observatory, Satavropol, Russia

Electronic mail: ntik@nik.sao.ru

YONG-IK BYUN

Department of Astronomy and Center for Space Astrophysics, Yonsei University, Seoul 120-749, Korea

Electronic mail: byun@darkskey.yonsei.ac.kr

AND

EUNHYEUK KIM

Department of Astronomy, Seoul National University, Seoul 151-742, Korea

Electronic mail: ekim@astro.snu.ac.kr

*Draft version February 1, 2008*

## ABSTRACT

We present deep *BVI* CCD photometry of the stars in the dwarf galaxy DDO 210. The color-magnitude diagrams of DDO 210 show a well-defined red giant branch (RGB) and a blue plume. The tip of the RGB is found to be at  $I_{\text{TRGB}} = 20.95 \pm 0.10$  mag. From this the distance to DDO 210 is estimated to be  $d = 950 \pm 50$  kpc. The corresponding distance of DDO 210 to the center of the Local Group is 870 kpc, showing that it is a member of the Local Group. The mean metallicity of the red giant branch stars is estimated to be  $[\text{Fe}/\text{H}] = -1.9 \pm 0.1$  dex. Integrated magnitudes of DDO 210 within the Holmberg radius ( $r_H = 110$  arcsec = 505 pc) are derived to be  $M_B = -10.6 \pm 0.1$  mag and  $M_V = -10.9 \pm 0.1$  mag. *B* and *V* surface brightness profiles of DDO 210 are approximately consistent with an exponential law with scale lengths  $r_s(B) = 161$  pc and  $r_s(V) = 175$  pc. The brightest blue and red stars in DDO 210 (BSG and RSG) are found to be among the faintest in the nearby galaxies with young stellar populations:  $\langle M_V(3) \rangle_{\text{BSG}} = -3.47 \pm 0.11$  mag and  $\langle M_V(3) \rangle_{\text{RSG}} = -4.75 \pm 0.13$  mag. An enhancement of the star formation rate in the recent past (several hundred Myrs) is observed in the central region of DDO 210. The opposite trend is observed in the outer region of the galaxy, suggesting a possible two-component structure of the kind disk/halo found in spiral galaxies. The real nature of this two-component structure must, however, be confirmed with more detailed observations.

[To appear in the *Astronomical Journal* in August 1999]

*Subject headings:* galaxy: evolution — galaxies: irregular — galaxies: individual (DDO 210) — galaxies: stellar content — galaxies: photometry — Distance scale

## 1. INTRODUCTION

DDO 210 is a faint dwarf galaxy in Aquarius discovered by van den Bergh (1959). It was resolved in ground based photographic images so that it was sometimes considered to be located nearby and to be a member of the Local Group (Fisher & Tully 1975, 1979; Yahil et al. 1977). It has been known to be a dwarf irregular galaxy. Lo et al. (1993) obtained an HI mass of  $3 \times 10^6 M_\odot$  and a total virial mass of  $1.4 \times 10^7 M_\odot$  in DDO 210, calculated assuming a distance of 1.0 Mpc. Taylor et al. (1998) gave an upper limit for the CO emission of  $I_{\text{CO}} < 0.11$  K km s<sup>-1</sup>.

The distance to DDO 210 has never been measured reliably and is not yet well known. From the proximity of DDO 210 on the sky and in velocity to NGC 6822, Fisher & Tully (1979) assumed that both galaxies were at the same distance (0.7 Mpc). On the other hand, from *BV* CCD photometry of bright stars in DDO 210, Greggio et al. (1993) suggested that this galaxy may be located at

as far as 4 Mpc, placing it beyond the edge of the Local Group. van den Bergh (1994) estimated the distance to be 0.8 Mpc, from simple comparison of the DDO 210 and the Leo I color-magnitude diagrams.

In this paper we present a study of stellar populations in DDO 210 and the Local Group membership of this galaxy based on deep ground based *BVI* CCD photometry. A brief progress report of this study was given in Lee (1999), which is superseded by this paper. This paper is composed as follows. Section 2 describes the observations and data reduction; Section 3 investigates the morphological structure of DDO 210. Section 4 presents the color-magnitude diagrams of the galaxy; and Section 5 estimates the distance to DDO 210 and concludes on the Local Group membership of DDO 210. Section 6 presents the surface photometry of DDO 210, Section 7 investigates the property of the brightest blue and red stars in DDO 210, and Section 8 discusses the star formation history (SFH). Finally, Section 9 summarizes the primary results.

<sup>1</sup> Based on observations made with the 2.5 m Nordic Optical Telescope operated on the island of La Palma by NOT S.A. in the Spanish Observatorio del Roque de Los Muchachos of the Instituto de Astrofísica de Canarias.

## 2. OBSERVATIONS AND DATA REDUCTION

### 2.1. Observations and Data Reduction

*BVI* CCD images of DDO 210 were obtained during three different observing runs at the Palomar 1.5 m telescope, the University of Hawaii 2.2 m telescope, and the NOT 2.5 m telescope of Roque de los Muchachos Observatory at La Palma. Table 1 lists the journal of observations of DDO 210. A grey scale map of the *V*-band CCD image of DDO 210 obtained at UH 2.2 m is displayed in Fig. 1 which shows that there are many bright foreground stars in the direction of this galaxy. For the analysis of the data, we have divided the field covered by our CCD images into three regions as shown in Fig. 1: the C-region covers the central main body of DDO 210, the I-region roughly extends to the boundaries of the optical body of the galaxy, the F-region is a control field and the F2-region represents the field outside the I-region in the Roque de los Muchachos field.

Instrumental magnitudes of the stars in the CCD images were derived using DoPHOT (Schechter et al. 1993) and DAOPHOT (Stetson 1987). Palomar and Roque de los Muchachos data were taken under photometric conditions and were calibrated using several standard stars (Landolt 1992) observed on the same runs. The calibration errors of each photometry are 0.04–0.05. The magnitudes of a subset of bright, well measured stars in common between the Palomar and Roque de los Muchachos photometries were averaged to obtain a common photometric scale to which all the magnitudes were transformed. The Hawaii data were also transformed to this photometric scale.

The Roque de los Muchachos *VI* photometry is the deepest one so that we have preferably used it for the analysis of the stellar populations in DDO 210. The Hawaii *BVI* data were used for the analysis of the bright stellar populations. The total number of stars which were measured in *V* and at least one other color with ALLSTAR  $\sigma$  values less than 0.25 is  $\sim 1800$ . Table 2 lists *BVI* photometry of the measured bright stars with  $V < 22$  mag inside the boundary of the I-region. *X* and *Y* coordinates increase to the east and to the south, respectively.

### 2.2. Comparison with Previous Photometry

Previous *BV* photometries of stars in DDO 210 were given by Marconi et al. (1990), Greggio et al. (1993, G93 from now on) and Hopp & Schulte-Ladbeck (1995, HP 95 from now on). The photometry by Marconi et al. (1990) was based on a wrong calibration and was corrected in G93. We have compared our photometry with these previous ones. The results of the comparison are  $\Delta V$  (this study – G93) =  $+0.01 \pm 0.09$ ,  $\Delta(B - V)$  (this study – G93) =  $+0.05 \pm 0.15$  for 30 common stars with  $V < 22$  mag, and  $\Delta V$  (this study – HP95) =  $-0.04 \pm 0.10$ ,  $\Delta(B - V)$  (this study – HP95) =  $+0.05 \pm 0.11$  for 35 common stars with  $V < 22$  mag. These results show a reasonable agreement in *V* and  $(B - V)$  among the three photometry sets.

## 3. MORPHOLOGICAL STRUCTURE

Fig. 1 shows that, morphologically, DDO 210 seems to have two components: the central component, represented by the C-region (of size  $2'.32 \times 0'.87$ ), and the extended component, represented by the I-region (of size  $4'.85 \times 2'.13$ ). From now on, by the I-region, we will refer

only to the outer part of this region, i.e., excluding the C-region. Both are elongated in the east-west direction, but have not common centers, the central component lying more east-ward than the extended one. A color-map created by combining *BVI* images shows that the stars in the central component are mostly very blue, while the stars in the outer region are mostly yellow to red. We have estimated the ellipticity and the position angle of the extended component to be roughly  $e = 0.5$  and  $PA = 173^\circ$ , respectively. Summarizing, DDO 210 is an elongated dwarf galaxy with a likely young stellar component close but out of its center. This confirms the common tendency of dwarf galaxies to show off-center star forming regions.

## 4. COLOR-MAGNITUDE DIAGRAMS

We display  $V - (B - V)$  and  $I - (V - I)$  diagrams of the measured stars in the C-region, I-region and F-region in Figs. 2 and 3. Note that the area of the (C+I)-region in the field is the same as that of the F-region so that we can estimate the contamination due to foreground stars by comparing the diagrams of each region.

Several distinguishable features of the stars in DDO 210 are seen in Figs. 2 and 3. First, there is a blue plume of bright stars with  $(B - V) < 0.4$  in the (C+I) region. Comparison of the (C+I)-region and the F-region in Fig. 2 shows that these bright blue stars are mostly members of DDO 210. The brightest end of the blue plume extends up to  $V \approx 21.0$  mag and  $(B - V) \approx -0.2$ . These stars are young stars and are mostly located in the C-region.

Secondly, there are several bright red stars with  $20 < V < 21$  mag and  $1.0 < (B - V) < 1.5$  in the C and I-region in Fig. 2. They might be young, red supergiants in DDO 210. However, the fact that a large number of stars is found in the F-region with  $V < 22$  mag (for every color); the fact that they are more numerous in the I-region than in the C-region, as it should be expected from the bigger size of field; and, more important, the fact that the blue population (which must be expected as a counterpart of the bright, red one) in the I-region is less numerous than in the C-region, would drive to the conclusion that many of the bright, red stars in field I are foreground.

Thirdly, Fig. 3 shows that there is a strong concentration of red stars fainter than  $I \approx 21$  mag and redder than  $(V - I) \approx 0.8$  mag in the C and I-regions. This feature corresponds to the red giant branch (RGB) and asymptotic giant branch (AGB) loci, populated by old and intermediate-age stars of DDO 210. The tip of the RGB (TRGB) is seen at  $(V - I) \approx 1.4$ .

## 5. DISTANCE AND METALLICITY

We estimate the distance to DDO 210 using the *I* magnitude of the TRGB, as described in Da Costa & Armandroff (1990) and Lee et al. (1993). The *I* magnitude of the TRGB is estimated using the  $I - (V - I)$  diagram in Fig. 3 and the luminosity function of red stars in the range defined by  $(V - I, I) = (0.5, 23.5)$ ,  $(1.2, 23.5)$  and  $(1.8, 19)$ . Fig. 4 shows the *I*-band luminosity function of the measured red stars in the (C+I)-region. We also plot the luminosity functions of stars of the same color range in the F-region and the F2-region in Fig. 4. This figure shows that while some star of DDO 210 can be still populating the F2-region, the F-region is very likely free of them. In

any case, for the purposes of this section, the contribution due to foreground stars is negligible for the bright part of the luminosity function of the (C+I) region. The luminosity function shows a sudden increase at  $I = 20.95 \pm 0.10$  mag. This corresponds to the TRGB seen in the color-magnitude diagram in Fig. 3. We will adopt this value as the  $I$  magnitude of the TRGB.

The mean color of the TRGB is estimated to be  $(V - I) = 1.44 \pm 0.04$ . The bolometric magnitude of the TRGB is then calculated from  $M_{\text{bol}} = -0.19[\text{Fe}/\text{H}] - 3.81$ . Adopting a mean metallicity of  $[\text{Fe}/\text{H}] = -1.86 \pm 0.12$  dex as estimated below, we obtain a value for the bolometric magnitude of  $M_{\text{bol}} = -3.46$  mag. From the equation  $\text{BC}_I = 0.881 - 0.243(V - I)_{\text{TRGB}}$  the bolometric correction at  $I$  for the TRGB is estimated to be  $\text{BC}_I = 0.54$  mag. The intrinsic  $I$  magnitude of the TRGB is then  $M_I = M_{\text{bol}} - \text{BC}_I = -4.00$  mag.

We adopt in this study the foreground reddening value  $E(B - V) = 0.03$  mag for DDO 210 in Burstein & Heiles (1982) and Ratnatunga & Bahcall (1985), corresponding to  $A_I = 0.06$  mag. This value is also similar to that given by Schlegel et al. (1998),  $E(B - V) = 0.05$ . Finally the distance modulus of DDO 210 is obtained:  $(m - M)_0 = 24.89 \pm 0.11$  mag, corresponding to a distance of  $950 \pm 50$  kpc.

The metallicity of an old stellar population can be estimated from the color of the RGB. DDO 210 has a composite stellar population containing also young stars, but an estimate of the metallicity using that approach is still quite useful, since no previous determinations exist. We have estimated the mean metallicity using the  $(V - I)$  color of the RGB stars 0.5 mag fainter than the TRGB,  $(V - I)_{-3.5}$ . This color is measured from the median value of the colors of red giant branch stars with  $I = 21.45 \pm 0.05$  mag in the I-region, resulting to be  $(V - I)_{-3.5} = 1.34 \pm 0.04$ . From this value we estimate the mean metallicity to be  $[\text{Fe}/\text{H}] = -1.86 \pm 0.12$  dex. This value is similar to those of dwarf irregular galaxies Sextans A ( $[\text{Fe}/\text{H}] = -1.9 \pm 0.1$  dex) and UKS 2323-326 ( $[\text{Fe}/\text{H}] = -1.9 \pm 0.1$  dex), and is close to the lowest end in the metallicity of stellar populations in dwarf irregular galaxies (Mateo 1998, Lee & Byun 1999).

In Fig. 5 we overlayed the loci of the red giant branches of Galactic globular clusters, of M15, M2, and NGC 1851, shifted according to the distance and reddening of DDO 210. The metallicities of M15, M2 and NGC 1851 are  $[\text{Fe}/\text{H}] = -2.17, -1.58$  and  $-1.29$  dex, respectively. Fig. 5 shows that the bright part of the RGB of DDO 210 is located well between those of M15 and M2. The broadening of the faint part of the RGB is mostly due to the photometric errors.

Greggio et al. (1993) estimated the distance to DDO 210 to be 4 Mpc, which is four times larger than our result. They derived this value from the comparison of their  $BV$  photometry of DDO 210 with the synthetic color-magnitude diagrams created using the Padova models of stellar evolution. They stated in their paper that “the true distance modulus of DDO 210 is  $(m - M)_0 \approx 28$ , although the uncertainty is obviously large”. If DDO 210 were at the distance of 4 Mpc, the absolute magnitude of the bright tip of the RGB would be  $M_I \approx -7$  mag. This is unrealistic, because the  $I$ -band absolute magnitude of the TRGB is known to be almost constant at  $M_I \approx -4.0$  mag for various types of resolved galaxies with old red giant

populations (Lee et al. 1993).

With the distance estimate obtained as above, we can tell whether DDO 210 is a member of the Local Group or not. Assuming that the center of the Local Group is, from our Galaxy, at two thirds of the connection line between our Galaxy and M31, we derive the distance to DDO 210 from the center of the Local Group to be 870 kpc. In addition, the systemic velocity of DDO 210 is  $v = -137 \pm 5$  km s $^{-1}$  (RC3), and the velocity with respect to the center of the Local Group is derived to be  $-15$  km s $^{-1}$ . From these values we conclude that DDO 210 is definitely a member of the Local Group. DDO 210 does not belong to either of the Milky Way subgroup or the M31 subgroup.

## 6. SURFACE PHOTOMETRY

The low surface brightness of DDO 210 and the presence of several bright foreground stars in the field, makes it difficult to derive reliably the surface photometry of the galaxy. We have proceeded for the surface photometry as follows. First, we removed several bright stars which are considered obviously to be foreground stars both from the DDO 210 and the F-region images using IMEDIT in IRAF. The mean intensity of the F-region was used for background sky estimation. Then we performed aperture photometry of DDO 210 using elliptical annular apertures with the ellipticity and position angle determined in Section 3. The results are listed in Table 3 and are displayed in Fig. 6. In Table 3  $r_{\text{eff}}$  represents a mean radius of the major axis of an ellipse, and  $r_{\text{out}}$  presents the outer radius of an elliptical aperture. Surface brightnesses are given in terms of mag arcsec $^{-2}$ .

Fig. 6 shows that the surface brightness profiles of DDO 210 follow approximately the exponential disk law. The scale lengths of DDO 210 are determined from these profiles to be  $r_s(B) = 35$  arcsec = 161 pc and  $r_s(V) = 38$  arcsec = 175 pc. The steep increase of the brightness in the center is due to a bright star located close to the center of the galaxy. The magnitude and color of this star (ID 1407) are  $V = 20.87 \pm 0.10$  and  $(B - V) = 1.14 \pm 0.10$ . With this color, it could be a main sequence star in the halo of the Milky Way, 5 kpc from the Sun or 7.5 kpc from the galactic center. It is also possible, however, that it is a yellow supergiant in DDO 210. The color profile of DDO 210 in Fig. 6 shows that the colors are almost constant around  $(B - V) \approx 0.25$  within  $r \approx 50$  arcsec (except for the central region), and get redder beyond  $r \approx 50$  arcsec. This shows that blue stellar populations are located mostly within  $r \approx 50$  arcsec.

The Holmberg radius corresponding to  $\mu_B = 26.5$  mag arcsec $^{-2}$  is approximately  $r_H = 110$  arcsec = 505 pc. The integrated magnitudes of DDO 210 within this radius are:  $B = 14.37 \pm 0.03$  mag,  $V = 14.06 \pm 0.03$  mag and  $I = 13.03 \pm 0.04$  mag. de Vaucouleurs et al. (1991, RC3) lists, as total magnitudes of DDO 210,  $B = 14.00 \pm 0.52$  mag and  $V = 13.88 \pm 0.52$  mag. These magnitudes are similar to ours, but the errors in our photometry are much smaller. The corresponding absolute magnitudes are  $M_B = -10.6 \pm 0.1$  mag,  $M_V = -10.9 \pm 0.1$  mag, and  $M_I = -11.9 \pm 0.1$  mag. This result shows that DDO 210 is the faintest among the known dwarf irregular galaxies in the Local Group (Mateo 1998).

## 7. THE BRIGHTEST BLUE AND RED STARS IN DDO 210

Little is known about the nature of the brightest stars in DDO 210. In the study of M supergiants in Local Group irregular galaxies Elias & Frogel (1985) stated “Certainly DDO 210 does not have a population of red supergiants comparable to that in Sextans A, which implies that it either is more distant or has no red supergiants.”. Since it turns out that DDO 210 is closer than Sextans A (this study, Mateo 1998), the only possibility is that there may be no red supergiants in DDO 210. Our photometry allow us to investigate the nature of the brightest stars in DDO 210.

We have determined the mean colors and magnitudes of the brightest stars. In the  $V - (B - V)$  diagram of the (C+I) region of DDO 210 shown in Fig. 2 it is obvious which stars are the three brightest blue stars in DDO 210 (IDs: 1656, 1503, and 1517), but it is not obvious which stars are the three brightest red stars because of contamination due to some foreground stars. We chose, as the three brightest red stars, three among the ones with  $20 < V < 21$  and  $1.0 < (B - V) < 1.5$  (IDs: 1468, 1071, and 926). The mean magnitudes and colors of these three brightest blue and red stars (called as BSG and RSG hereafter) in DDO 210 are derived to be, respectively,  $\langle V(3) \rangle_{BSG} = 21.51 \pm 0.11$  mag and  $\langle (B - V)(3) \rangle_{BSG} = -0.15 \pm 0.03$ , and  $\langle V(3) \rangle_{RSG} = 20.23 \pm 0.13$  mag and  $\langle (B - V)(3) \rangle_{RSG} = 1.13 \pm 0.12$ . The corresponding absolute magnitudes and colors are  $\langle M_V(3) \rangle_{BSG} = -3.47$  mag,  $\langle (B - V)(3) \rangle_{BSG} = -0.18$ ,  $\langle M_V(3) \rangle_{RSG} = -4.75$  mag and  $\langle (B - V)(3) \rangle_{RSG} = 1.10$ , respectively. If some of the chosen brightest red stars are foreground stars, the mean magnitudes of the brightest red stars in DDO 210 will be fainter.

We have compared the magnitudes and colors of the brightest stars in DDO 210 with those of other galaxies with young stellar populations in Fig. 7. The data for other galaxies are from Lyo & Lee (1997). Fig. 7 shows that both the brightest blue and red stars in DDO 210 are the faintest among the sample galaxies. This result is consistent with the fact that DDO 210 is the faintest among the known dwarf irregular galaxies in the Local Group. Fig. 7 shows also that the brightest red stars are the least red among the sample galaxies. This can be interpreted in the sense that the brightest red stars in DDO 210 may be more metal-poor than those in brighter galaxies.

## 8. THE STAR FORMATION HISTORY OF DDO210

From a qualitative point of view, the presence of a conspicuous RGB-AGB structure implies star formation at intermediate to old ages in DDO 210. Moreover, the age of the youngest population in DDO 210 can be estimated from comparison of the CMD of Fig. 2 with theoretical isochrones. Fig. 8 displays the  $V - (B - V)$  CMD of DDO 210 together with three isochrones for metallicities  $Z = 0.0004$  and ages 30, 100 and 300 Myrs from the Padova library (see Bertelli et al. 1994 and references therein). It shows that star formation as recent as some 30 Myrs ago has taken place in the galaxy.]

The star formation history (SFH) can be derived quantitatively in detail from deep CMDs as shown by Gallart et al. (1999). The CMD of DDO 210 is not deep enough for such a derivation, but it can still be used to sketch the

SFH. For this purpose, stars with  $-4 < M_{I,0} < -2.5$  have been counted in the CMD of fields C-region, I-region and F2-region and divided into two groups: bluer and redder than  $(V - I)_0 = 0.5$ . After normalization to the same area, the counts of the F2-region have been subtracted from those of the C and I-regions to correct of foreground contamination. The relatively strong foreground contamination of the DDO 210 field prevents us from using the stars over the TRGB for the calculation of the SFH. On the other hand, the maximum magnitude used ( $M_I = -2.5$  mag) and the fact that we pursue only a rough estimate of the SFR, allow us to neglect crowding effects.

Following a method similar to that described in Aparicio et al. (1997b) and Aparicio et al. (1999), a synthetic CMD has been calculated with 100,000 stars brighter than  $M_I = -1.5$  mag, a constant SFR and random metallicities in the interval  $0.0001 < Z < 0.0003$ , which corresponds to the interval of  $[\text{Fe}/\text{H}]$  found from the RGB of DDO 210. The synthetic stars have been divided into two age intervals: younger and older than 1 Gyr, and the same boxes as for the observational CMDs have been defined in each of the resulting synthetic CMDs. The resulting SFRs for the 15 to 1 Gyrs ago and the 1 to 0 Gyrs ago are given in Table 4 and shown in Fig. 9 for the central (C) and the outer (I) regions of DDO 210. These values are also given after surface normalization (right-hand ordinate axis of Fig. 9). The areas used for those normalizations have been  $2.18 \text{ arcsec}^2$  or  $1.67 \times 10^5 \text{ pc}^2$  for the C-region and  $8.11 \text{ arcsec}^2$  or  $6.21 \times 10^5 \text{ pc}^2$  for the I-region.

DDO 210 shows a SFH behavior in its central region similar to what has been found in other dIrr galaxies, like NGC 6822 (Gallart et al. 1996), Pegasus (Aparicio et al. 1997a), Antlia (Aparicio et al. 1997c) and DDO 187 (Aparicio et al. 1999). In overall, it consists in an enhancement of the SFR for the recent past of the history of the galaxy with respect to the value averaged for its entire life. This can be interpreted as the result of a fluctuating star formation rate (see Aparicio et al. 1999). It is also interesting to note that this recent strength of the SFR does not affect the entire galaxy but the central part only. For the outer I-region, the SFR decreased for recent epochs. This behaviour is common to other dIrr: WLM (Minniti, & Zijlstra 1996), Antlia (Aparicio et al. 1997c), Phoenix (Martínez-Delgado et al. 1999), DDO 187 (Aparicio et al. 1999) and may suggest a two-components structure of the kind disk-halo of bigger spiral galaxies. However, more detailed data are needed to firmly state such conclusion.

## 9. SUMMARY AND CONCLUSION

We have presented a study of the stellar populations and the Local Group membership of the dwarf galaxy DDO 210 based on deep *BVI* CCD photometry. The primary results obtained in this study are summarized as follows.

(1) *BVI* color-magnitude diagrams of  $\sim 1800$  stars in the  $7'.5 \times 7'.5$  area of DDO 210 have been presented. These color-magnitude diagrams exhibit a well-defined RGB and a blue plume.

(2) The TRGB is found to be at  $I = 20.95 \pm 0.10$  mag and  $(V - I) = 1.44 \pm 0.04$  mag. From this, a distance modulus of  $(m - M)_0 = 24.89 \pm 0.11$  mag is derived, corresponding to a distance of  $950 \pm 50$  kpc. The distance to the barycenter of the Local Group is derived to be 870 kpc. From this result and the systemic velocity of DDO

210 we conclude that DDO 210 is definitely a member of the Local Group.

(3) The mean color of the RGB at  $M_I = -3.5$  mag is  $(V - I) = 1.34 \pm 0.04$  mag. From this value we obtain a mean metallicity of the RGB:  $[\text{Fe}/\text{H}] = -1.86 \pm 0.12$  dex.

(4)  $B$  and  $V$  surface brightness profiles of DDO 210 follow roughly the exponential disk law with scale lengths of  $r_s(B) = 35$  arcsec = 161 pc and  $r_s(V) = 38$  arcsec = 175 pc. The Holmberg radius corresponding to  $\mu_B = 26.5$  mag arcsec $^{-2}$  is  $r_H = 110$  arcsec = 505 pc. The integrated magnitudes up to this radius are  $M_B = -10.6$  mag,  $M_V = -10.9$  mag, and  $M_I = -11.9$  mag.

(5) The magnitudes of the brightest blue and red stars in DDO 210 are derived:  $\langle M_V(3) \rangle_{BSG} = -3.47 \pm 0.11$  mag and  $\langle M_V(3) \rangle_{RSG} = -4.75 \pm 0.13$  mag. The brightest blue and red stars in DDO 210 are found to be among the faintest in nearby galaxies with young stellar populations.

(6) The SFRs of the central and extended regions of

DDO 210 have been calculated. The central region shows an enhancement of the star formation activity in recent epochs, in agreement to what has been observed in other dwarf irregular galaxies. In the extended region of DDO 210, the SFR has been lower in the recent past than the value averaged for the entire life of the galaxy, indicating a possible two-components structure for the galaxy that must, however, be confirmed with more detailed data.

This research is supported by the Ministry of Education, Basic Science Research Institute grant No.BSRI-98-5411 (to M.G.L.); by the Instituto de Astrofísica de Canarias, grant PB3/94 and the DGES of the Kingdom of Spain, grant PB97-1438-C02-01 (to A.A.); by Creative Research Initiatives Program of the Korean Ministry of Science and Technology and also by Yonsei University Research Grant (to Y.I.B.).

#### REFERENCES

- Aparicio, A., Gallart, C., & Bertelli, G. 1997a, *AJ*, 114, 669  
Aparicio, A., Gallart, C., & Bertelli, G. 1997b, *AJ*, 114, 680  
Aparicio, A., Dalcanton, J., Gallart, C., & Martínez-Delgado, D. 1997c, *AJ*, 114, 1447  
Aparicio, A., Tikonov, N., & Karachentsev, I. 1999, *AJ*, submitted  
Bertelli, G., Bressan, A., Fagotto, F., Chiosi, C., & Nasi, E. 1994, *A&AS*, 106, 275  
Burstein, D., & Heiles, C. 1982, *AJ*, 87, 65  
Da Costa, G. S., & Armandroff, T. E. 1990, *AJ*, 100, 162  
de Vaucouleurs, G., de Vaucouleurs, A., Corwin Jr., H. G., Buta, R. J., Paturel, G., and Fouque, P. 1991, *Third Reference Catalog of Bright Galaxies* (Springer-Verlag, New York) (RC3)  
Elias, J. K., & Frogel, J. A. 1985, *AJ*, 289, 141  
Fisher, J. R., & Tully, R. B. 1975, *A&A*, 44, 151  
Fisher, J. R., & Tully, R. B. 1979, *AJ*, 84, 62  
Gallart, C., Aparicio, A., Bertelli, G., & Chiosi, C. 1996, *AJ*, 112, 1950  
Gallart, C., Freedman, W.L., Aparicio, A., Bertelli, G., & Chiosi, C. 1999, *AJ*, submitted  
Greggio, L., Marconi, G., Tosi, M., & Focardi, P. 1993, *AJ*, 105, 894  
Hopp, U., & Schulte-Ladbeck, R. E. 1995, *A&A*, 111, 527  
Landolt, A. U. 1992, *AJ*, 104, 340  
Lee, M. G., Freedman, W. L., & Madore, B. F. 1993, *ApJ*, 417, 553  
Lee, M. G. 1999, *IAU Symp. No. 192, The Stellar Content of the Local Group Galaxies*, in press  
Lee, M. G., & Byun, Y.-I. 1999, *AJ*, in press (August issue)  
Lo, K. Y., Sargent, W. L. W., & Young, K. 1993, *AJ*, 106, 507  
Lyo, A.-R., & Lee, M. G. 1997, *JKAS*, 30, 27  
Marconi, G., Focardi, P., Greggio, L., & Tosi, M. 1990, *ApJ*, 360, 39L  
Martínez-Delgado, D., Gallart, C., & Aparicio, A. 1999, *AJ*, submitted  
Mateo, M. 1998, *ARA&A*, 36, 435  
Minniti, D., & Zijlstra, A. A. 1996, *ApJ*, 463, L13  
Ratnatunga, K. U., & Bahcall, J. N. 1985, *ApJS*, 59, 63  
Schechter, P., Saha, A., & Mateo, M. L. 1993, *PASP*, 105, 1342  
Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525  
Stetson, P. B. 1987, *PASP*, 99, 191  
Taylor, C. L., Kobulnicky, H. A., & Skillman, E. D. 1998, *AJ*, 116, 2746  
van den Bergh, S. 1959, *Publ. David Dunlop Obs. II. No. 5*  
van den Bergh, S. 1994, *AJ*, 107, 1328  
Yahil, A., Tammann, G. A., & Sandage, A. 1977, *ApJ*, 217, 903




FIG. 1.— A grey-scale map of the V-band CCD image of DDO 210. North is at the top and east is to the left. The size of the field is  $7'.5 \times 7'.5$ . Rectangular regions labeled as C, I, and F represent respectively, the central region, the intermediate region and the control field region. The square labeled F2 in the lower middle area represents the field covered by the Roque de los Muchachos observations.




FIG. 2.—  $V-(B-V)$  diagram of the measured stars in the C-region (a), the I-region (b) and the F-region (c) of DDO 210.




FIG. 3.— (a)  $I-(V-I)$  diagram of the measured stars in the C-region (a), the I-region (b) and the F-region (c) of DDO 210. Open squares in (c) represent the stars in the F2-region.






FIG. 4.—  $I$ -band luminosity function of the red giant branch stars in the C region plus I-region. The tip of the red giant branch is labeled as TRGB. The solid line and dotted lines represent the luminosity functions of the F-region and F2-region, respectively.




FIG. 5.—  $I$ – $(V - I)$  diagram of the measured stars in the I-region (filled circles) and C-region (open circles) of DDO 210 in comparison with the red giant branches of Galactic globular clusters. The solid curved lines show, from left to right, the loci of the giant branches of M15, M2, and NGC 1851, the metallicities of which are  $[\text{Fe}/\text{H}] = -2.17$ ,  $-1.58$  and  $-1.29$  dex, respectively. The mean errors for the magnitudes and colors are illustrated by the error bars at the right.




FIG. 6.— Surface photometry of DDO 210. (a) Surface brightness profiles vs. effective radius along the major axis.  $B$ ,  $V$  and  $I$  magnitudes are represented by the filled circles, open circles and crosses. The solid line represents a linear fit to the  $B$  data. (b) Differential colors vs effective radius along the major axis.




FIG. 7.—  $V-(B-V)$  diagram of the brightest blue and red supergiants in DDO 210 (filled squares) in comparison with other galaxies (open squares). The magnitudes and colors are average values of the three brightest blue and red supergiants in each sample galaxy.




FIG. 8.—  $V-(B-V)$  diagram of the measured stars in the C-region (open circles) of DDO 210 in comparison with theoretical isochrones. The solid lines represent the Padova isochrones for the metallicity of  $Z = 0.0004$  and ages of 30, 100, and 300 Myrs. The mean errors for the magnitudes and colors are illustrated by the error bars at the right.




FIG. 9.— The star formation history of DDO 210. The bottom part of the right-hand ordinate axis scale refers to the surface normalized  $\psi(t)$  of the external region of the galaxy. The upper part of the same axis refers to the central region. The values of  $\psi(t)$  for the two age intervals plotted (15–1 and 1–0 Gyr ago) must be considered as the averages for those intervals. The error bars correspond to poissonian statistics on the star counts. The dashed lines show the intervals of time over which the averages have been done, but are not intended to give accurate information about the detailed shape of the star formation history.

TABLE 1  
JOURNAL OF OBSERVATIONS FOR DDO 210

| Filters  | T <sub>exp</sub> | Air mass | FWHM  | U.T.(Start)       | Telescope     |
|----------|------------------|----------|-------|-------------------|---------------|
| <i>B</i> | 1200 s           | 1.56     | 1''.2 | 1992 Aug 23 07:52 | Palomar 1.5 m |
| <i>V</i> | 900 s            | 1.65     | 1''.0 | 1992 Aug 23 08:12 |               |
| <i>I</i> | 600 s            | 1.45     | 0''.9 | 1992 Aug 23 06:25 |               |
| <i>B</i> | 1000 s           | 1.33     | 1''.2 | 1994 Oct 9 07:32  | UH 2.2 m      |
| <i>V</i> | 3 × 900 s        | 1.20     | 1''.0 | 1994 Oct 9 06:05  |               |
| <i>I</i> | 3 × 600 s        | 1.25     | 0''.9 | 1994 Oct 9 06:50  | NOT 2.5 m     |
| <i>V</i> | 3 × 900 s        | 1.21     | 0''.8 | 1997 Jul 27 00:36 |               |
| <i>I</i> | 4 × 600 s        | 1.26     | 0''.9 | 1997 Jul 27 01:10 |               |

TABLE 2.  
PHOTOMETRY OF THE BRIGHT STARS WITH  $V < 22$  MAG IN DDO 210

| ID   | X(pixel) | Y(pixel) | <i>V</i> | ( <i>B</i> − <i>V</i> ) | ( <i>V</i> − <i>I</i> ) | ID   | X(pixel) | Y(pixel) | <i>V</i> | ( <i>B</i> − <i>V</i> ) | ( <i>V</i> − <i>I</i> ) |
|------|----------|----------|----------|-------------------------|-------------------------|------|----------|----------|----------|-------------------------|-------------------------|
| 1782 | 526.3    | 789.4    | 16.85    | 0.61                    | 0.83                    | 1407 | 462.0    | 702.5    | 20.93    | 1.14                    | 1.35                    |
| 1029 | 405.3    | 641.8    | 17.09    | 0.88                    | 1.06                    | 1829 | 118.1    | 814.4    | 20.96    | 1.29                    | 2.24                    |
| 1755 | 358.8    | 778.7    | 17.19    | 0.50                    | 0.87                    | 1362 | 317.2    | 694.8    | 21.01    | 0.18                    | 0.22                    |
| 1149 | 300.1    | 661.4    | 17.48    | 0.64                    | 0.84                    | 1656 | 487.9    | 745.8    | 21.05    | -0.17                   | 0.04                    |
| 1571 | 538.4    | 726.6    | 17.88    | 0.55                    | 0.38                    | 1710 | 468.5    | 761.3    | 21.05    | 0.23                    | 0.63                    |
| 1423 | 461.5    | 704.5    | 18.12    | 0.86                    | 1.02                    | 1189 | 668.9    | 667.5    | 21.20    | 1.03                    | 0.85                    |
| 1763 | 598.6    | 783.2    | 18.35    | 0.90                    | 1.35                    | 1548 | 441.5    | 722.1    | 21.24    | 0.57                    | 0.61                    |
| 1202 | 668.3    | 669.4    | 18.45    | 0.90                    | 1.13                    | 799  | 176.6    | 558.5    | 21.30    | 1.07                    | 1.67                    |
| 855  | 611.0    | 590.1    | 19.32    | 0.59                    | 0.73                    | 910  | 675.6    | 620.1    | 21.39    | 1.12                    | 1.71                    |
| 918  | 668.2    | 622.5    | 19.72    | 0.64                    | 0.91                    | 1477 | 461.5    | 709.6    | 21.40    | 0.41                    | 0.72                    |
| 1243 | 323.0    | 675.7    | 19.73    | 1.50                    | 2.23                    | 1506 | 470.3    | 714.7    | 21.44    | 0.91                    | 1.15                    |
| 1780 | 287.7    | 788.0    | 19.84    | 0.42                    | 0.77                    | 820  | 291.7    | 571.0    | 21.45    | 1.01                    | 1.48                    |
| 1468 | 490.5    | 708.8    | 20.14    | 1.13                    | 1.74                    | 1503 | 464.4    | 713.9    | 21.48    | -0.15                   | 0.19                    |
| 1805 | 303.0    | 801.4    | 20.29    | 0.57                    | 0.94                    | 1632 | 515.3    | 739.2    | 21.50    | 0.12                    | 0.62                    |
| 1071 | 685.8    | 647.1    | 20.32    | 1.25                    | 2.14                    | 1517 | 523.8    | 717.2    | 21.54    | -0.14                   | 0.34                    |
| 926  | 425.1    | 625.0    | 20.40    | 1.01                    | 1.64                    | 1713 | 321.5    | 761.6    | 21.59    | 0.77                    | 1.38                    |
| 1444 | 231.0    | 706.2    | 20.40    | 0.27                    | 0.73                    | 1524 | 560.7    | 718.2    | 21.69    | -0.14                   | -0.10                   |
| 1830 | 255.9    | 815.4    | 20.57    | 0.70                    | 1.22                    | 1339 | 563.5    | 690.8    | 21.72    | 0.92                    | 0.49                    |
| 1821 | 252.7    | 811.5    | 20.70    | 0.84                    | 1.27                    | 1337 | 410.0    | 690.5    | 21.90    | 1.13                    | 1.62                    |
| 1206 | 171.2    | 669.8    | 20.75    | 1.35                    | 2.24                    | 1353 | 452.4    | 693.3    | 21.95    | 0.07                    | 1.28                    |
| 1600 | 366.1    | 733.0    | 20.77    | 0.32                    | 0.89                    | 1831 | 424.8    | 815.9    | 21.96    | 1.02                    | 1.51                    |
| 1551 | 368.7    | 722.3    | 20.77    | 1.30                    | 2.36                    | 1344 | 420.6    | 691.7    | 21.99    | 0.56                    | 0.61                    |

TABLE 3  
BVI SURFACE PHOTOMETRY OF DDO 210

| $R_{\text{eff}} [']$ | $\mu_B$          | $\mu_V$          | $\mu_I$          | $R_{\text{out}} [']$ | $B$   | $V$   | $I$   |
|----------------------|------------------|------------------|------------------|----------------------|-------|-------|-------|
| 6.2                  | $22.74 \pm 0.43$ | $22.31 \pm 0.53$ | $21.26 \pm 0.68$ | 8.8                  | 17.54 | 17.08 | 16.02 |
| 13.9                 | $23.74 \pm 0.19$ | $23.50 \pm 0.23$ | $22.55 \pm 0.44$ | 17.6                 | 16.69 | 16.34 | 15.32 |
| 22.4                 | $23.78 \pm 0.31$ | $23.59 \pm 0.25$ | $22.89 \pm 0.23$ | 26.4                 | 16.01 | 15.73 | 14.83 |
| 31.1                 | $24.16 \pm 0.26$ | $23.96 \pm 0.22$ | $23.17 \pm 0.17$ | 35.2                 | 15.59 | 15.34 | 14.47 |
| 39.8                 | $24.25 \pm 0.27$ | $24.06 \pm 0.24$ | $23.22 \pm 0.18$ | 44.0                 | 15.23 | 15.00 | 14.14 |
| 48.6                 | $24.40 \pm 0.34$ | $24.21 \pm 0.31$ | $23.20 \pm 0.37$ | 52.8                 | 14.95 | 14.73 | 13.83 |
| 57.4                 | $24.77 \pm 0.33$ | $24.55 \pm 0.29$ | $23.62 \pm 0.32$ | 61.6                 | 14.75 | 14.53 | 13.63 |
| 66.1                 | $25.01 \pm 0.16$ | $24.67 \pm 0.19$ | $23.38 \pm 0.41$ | 70.4                 | 14.61 | 14.37 | 13.40 |
| 74.9                 | $25.55 \pm 0.24$ | $25.20 \pm 0.28$ | $24.04 \pm 0.58$ | 79.2                 | 14.52 | 14.27 | 13.29 |
| 83.7                 | $25.82 \pm 0.28$ | $25.47 \pm 0.38$ | $24.11 \pm 0.53$ | 88.0                 | 14.44 | 14.19 | 13.20 |
| 92.5                 | $25.92 \pm 0.33$ | $25.43 \pm 0.40$ | $23.77 \pm 0.60$ | 96.8                 | 14.38 | 14.11 | 13.09 |
| 101.3                | $26.34 \pm 0.40$ | $26.03 \pm 0.53$ | $24.82 \pm 1.03$ | 105.6                | 14.33 | 14.06 | 13.03 |
| 110.1                | $26.49 \pm 0.38$ | $26.23 \pm 0.60$ | $24.29 \pm 0.60$ | 114.4                | 14.29 | 14.02 | 12.99 |
| 118.9                | $26.27 \pm 0.49$ | $25.81 \pm 0.48$ | $24.26 \pm 0.51$ | 123.2                | 14.23 | 13.97 | 12.92 |
| 127.7                | $26.46 \pm 0.59$ | $26.03 \pm 0.48$ | $24.52 \pm 0.62$ | 132.0                | 14.19 | 13.92 | 12.86 |



TABLE 4  
STAR FORMATION RATES

|                                                                                    | DDO 210 center | DDO 210 outer |
|------------------------------------------------------------------------------------|----------------|---------------|
| $\bar{\psi}_{15-1}$ ( $10^{-4} \text{ M}_{\odot}\text{yr}^{-1}$ )                  | 0.44           | 0.29          |
| $\bar{\psi}_{1-0}$ ( $10^{-4} \text{ M}_{\odot}\text{yr}^{-1}$ )                   | 1.14           | 0.09          |
| $\bar{\psi}_{15-1}/A$ ( $10^{-10} \text{ M}_{\odot}\text{yr}^{-1}\text{pc}^{-2}$ ) | 2.66           | 0.47          |
| $\bar{\psi}_{1-0}/A$ ( $10^{-10} \text{ M}_{\odot}\text{yr}^{-1}\text{pc}^{-2}$ )  | 6.87           | 0.15          |